

DIFFERENTIAL SUBSTITUTIONS AND SYMMETRIES OF HYPERBOLIC EQUATIONS

S. Ya. Startsev

Mathematical Institute of Ufa Center of Russian Academy,
Chernyshevsky Street 112, 450000 Ufa, Russia
e-mail: starts@nkc.bashkiriya.su

Abstract: There are considered differential substitutions of the form $v = P(x, u, u_x)$ for which there exists a differential operator $H = \sum_{i=0}^k \alpha_i D_x^i$ such that the differential substitution maps the equation $u_t = H[s(x, P, D_x(P), \dots, D_x^m(P))]$ into an evolution equation for any function s and any nonnegative integer m . All differential substitutions of the form $v = P(x, u, u_x)$ known to the author have this property. For example, the well-known Miura transformation $v = u_x - u^2$ maps any equation of the form

$$u_t = (D_x^2 + 2uD_x + 2u_x)[s(x, u_x - u^2, D_x(u_x - u^2), \dots, D_x^k(u_x - u^2))]$$

into the equation

$$v_t = (D_x^3 + 4vD_x + 2v_x)[s(x, v, \frac{\partial v}{\partial x}, \dots, \frac{\partial^k v}{\partial x^k})].$$

The complete classification of such differential substitutions is given. An infinite set of the pairwise nonequivalent differential substitutions with the property mentioned above is constructed. Moreover, a general result about symmetries and invariant functions of hyperbolic equations is obtained (see theorem 1).

Let us consider a differential substitution of the form

$$v = P(x, u, u_x) \tag{1}$$

which maps solutions of an evolution equation into solutions of another evolution equation. It is sufficient for our aim to give the following definition of the differential substitution (see [1] for more general definition).

Definition 1: (1) is called a differential substitution from an equation

$$u_t = f(x, u, u_1, \dots, u_n) \tag{2}$$

into an equation

$$v_t = g(x, v, v_1, \dots, v_n) \tag{3}$$

if $P_{u_x} \neq 0$ and the relationship

$$(P_{u_x} D_x + P_u)(f) = g(x, P, D_x(P), \dots, D_x^n(P))$$

is satisfied. Here u_i and v_i are the i -th partial derivatives of u and v with respect to x , respectively, D_x is the total derivative with respect to x expressed in the variables x, u, u_i as

$$D_x = \frac{\partial}{\partial x} + u_1 \frac{\partial}{\partial u} + \sum_{i=1}^{\infty} u_{i+1} \frac{\partial}{\partial u_i} .$$

It should be noted that differential substitutions are determined up to invertible (contact and point) transformations of the equations (2) and (3). A differential substitution π_1 is said to be *equivalent* to a differential substitution π_2 if $\pi_1 = \rho_2 \circ \pi_2 \circ \rho_1$, where ρ_1 and ρ_2 are invertible transformations of the equations (2) and (3), respectively. It is shown in [1] that any differential substitution

$$\begin{aligned} v &= \psi(x, u, u_1, u_2) \\ \bar{x} &= \varphi(x, u, u_1, u_2) \end{aligned}$$

which satisfies the conditions

$$\varphi_{u_2} D_x(\psi) = \psi_{u_2} D_x(\varphi), \quad \frac{\partial^2}{\partial u_2^2} \left(\frac{\varphi_u D_x(\psi) - \psi_u D_x(\varphi)}{\varphi_{u_1} D_x(\psi) - \psi_{u_1} D_x(\varphi)} \right) = 0 \quad (4)$$

is equivalent to a differential substitution of the form (1). Therefore, our study of the differential substitutions (1) is the study of the more wide class of differential substitutions. For example, all substitutions of the form

$$\begin{aligned} v &= \psi(x, u, u_1) \\ \bar{x} &= \varphi(x, u, u_1) \end{aligned}$$

satisfy the conditions (4).

We shall assume in this paper that for the differential substitution (1) there exists a differential operator

$$H = \sum_{i=0}^k \alpha_i D_x^i$$

where $k \geq 0$, α_i are functions of x, u, u_i and $\alpha_k \neq 0$, such that the differential substitution transforms the equation

$$u_t = H[s(x, P, D_x(P), \dots, D_x^m(P))]$$

into an evolution equation for any function s and any nonnegative integer m . All differential substitutions (1) known to the author have this property.

Examples: The example of the Miura transformation is contained in Abstract. In this case $H = D_x^2 + 2uD_x + u_x$.

Other examples are:

$$\begin{aligned} v &= u_1, & H &= 1; \\ v &= u_1 + e^u, & H &= D_x + u_1; \\ v &= u_1 + e^u + e^{-u}, & H &= D_x^2 + \frac{u_1 + e^{-2u} - e^{2u}}{u_1 + e^u + e^{-u}} D_x + \\ & & & + \frac{u_3 - 2u_1(e^{2u} + e^{-2u})}{u_1 + e^u + e^{-u}} - \left(\frac{u_2 + u_1(e^u - e^{-u})}{u_1 + e^u + e^{-u}} \right)^2 - u_1^2. \end{aligned}$$

It should be noted that these examples show very well that integrability of an equation and existence of a differential substitution relating this equation to another equation are not the same.

The present work is based on the fact that (1) is the differential substitution from the equation (2) into an evolution equation if and only if the right-hand side of the equation (2) is a symmetry of the hyperbolic equation

$$u_{xy} = -P_u/P_{u_1}u_y. \quad (5)$$

This fact was proved in [1].

We remind that a function f depending on $x, u, u_i, w_i = \frac{\partial^i u}{\partial y^i}$ is called a *symmetry* of the equation

$$u_{xy} = F(x, y, u, u_x, u_y) \quad (6)$$

if f satisfies the condition

$$(D_x D_y - F_{u_x} D_x - F_{u_y} D_y - F_u)(f) = 0,$$

where D_x and D_y are the total derivatives with respect to x and y by virtue of the equation (6), respectively. It should be noted that we eliminate the derivatives $\frac{\partial^{i+j} u}{\partial x^i \partial y^j}$ with $i \cdot j \neq 0$ by virtue of the equation (6) and its differential consequences. Therefore, we assume from now on that all functions depend on the variables x, y, u, u_i, w_i . The total derivatives D_x and D_y are expressed in these variables as

$$D_x = \frac{\partial}{\partial x} + u_1 \frac{\partial}{\partial u} + \sum_{i=1}^{\infty} \left(u_{i+1} \frac{\partial}{\partial u_i} + D_y^{i-1}(F) \frac{\partial}{\partial w_i} \right)$$

$$D_y = \frac{\partial}{\partial y} + w_1 \frac{\partial}{\partial u} + \sum_{i=1}^{\infty} \left(w_{i+1} \frac{\partial}{\partial w_i} + D_x^{i-1}(F) \frac{\partial}{\partial u_i} \right)$$

Definition 2: A function Q is said to be D_x -invariant (D_y -invariant) if $D_x(Q) = 0$ ($D_y(Q) = 0$). If Q depend on only y (only x), then Q is called a trivial D_x -invariant (D_y -invariant) function.

It is important to observe that any function of $x, P, D_x^i(P)$ is D_y -invariant for the equation (5) and the converse statement also is true. Using this fact, our assumption and the following theorem, we can reduce the classification problem for the differential substitutions (1) to the classification problem for the equations (5) which admit a nontrivial D_x -invariant function.

Theorem 1: Let an equation (6) admits a nontrivial D_y -invariant function and there exist a differential operator

$$H = \sum_{i=0}^k \alpha_i D_x^i$$

such that $H(s)$ is the symmetry of the equation (6) for any D_y -invariant function s . Then the equation (6) admits a nontrivial D_x -invariant function and there exist a differential operator

$$R = \sum_{i=0}^m \beta_i D_y^i$$

such that $R(q)$ is the symmetry of the equation (6) for any D_x -invariant function q .

Here α_i, β_i are functions of x, y, u and the derivatives of u ; $k, m \geq 0, \alpha_k, \beta_m \neq 0$.

The main idea of the proof is to demonstrate finiteness of the Laplace Y -index of the equation (6) (the definition of the Laplace indices is contained in [2]). The latter implies existence of the D_x -invariant function and the operator R (see [3]-[5]). The technique of the proof of finiteness of the Laplace Y -index is similar to one which is used in [2].

Let f be a symmetry of the equation (6). It is easy to verify that the vector field

$$\partial_f = f \frac{\partial}{\partial u} + \sum_{i=1}^{\infty} \left(D_y^i(f) \frac{\partial}{\partial w_i} + D_x^i(f) \frac{\partial}{\partial u_i} \right)$$

commutes with the total derivatives D_x and D_y . Hence, the operator

$$Q_* = \frac{\partial Q}{\partial u} + \sum_{i=1}^{\infty} \left(\frac{\partial Q}{\partial w_i} D_y^i + \frac{\partial Q}{\partial u_i} D_x^i \right)$$

maps any symmetry into a D_x -invariant (D_y -invariant) function if Q is D_x -invariant (D_y -invariant). Let us consider the equation (5). The straightforward calculation shows that $w_1 g$ is the symmetry of (5) for any D_x -invariant functions g . Therefore, if Q is D_x -invariant, then the operator $Q_* \circ w_1 = \sum_{i=0}^k c_i D_y^i$ maps any D_x -invariant function again into a D_x -invariant function and the coefficient c_i must be D_x -invariant. Taking this into account we can easily prove the following result which was obtained in a more complicated way in [6].

Proposition 1: *Let an equation*

$$u_{xy} = b(x, u, u_x) u_y$$

has a nontrivial D_x -invariant function. Then this equation can be reduced to another of the same form admitting the D_x -invariant function

$$Q = \frac{w_3}{w_1} - \frac{3}{2} \left(\frac{w_2}{w_1} \right)^2 \quad (7)$$

by a transformation

$$u = \psi(x, u). \quad (8)$$

The equation (5) has a nontrivial D_x -invariant function by our assumption and theorem 1. Conversely, if the equation (5) admits a nontrivial D_x -invariant function, then there exist a differential operator $H = \sum_{i=0}^k \alpha_i D_x^i$ which maps any D_y -invariant function of the equation (5) into the right-hand side of an evolution equation admitting the substitution (1) (see [4]). Transformations (8) correspond to point transformations of the equation (2). Thus differential substitutions (1) interesting for us are equivalent to ones which correspond to the equation (5) admitting the D_x -invariant function (7).

It is convenient for further consideration to express u_x from (1) and rewrite the differential substitution (1) as a system

$$\begin{cases} u_x = a(x, u, v) \\ v_y = 0 \end{cases} \quad (9)$$

We eliminate all partial derivatives of u with respect to x by virtue of (9). The total derivatives D_x and D_y are expressed in the remaining variables x, u, v, v_i, w_i as

$$\begin{aligned} D_x &= \frac{\partial}{\partial x} + a \frac{\partial}{\partial u} + v_1 \frac{\partial}{\partial v} + \sum_{i=1}^{\infty} \left(v_{i+1} \frac{\partial}{\partial v_i} + D_y^i(a) \frac{\partial}{\partial w_i} \right) \\ D_y &= \frac{\partial}{\partial y} + w_1 \frac{\partial}{\partial u} + \sum_{i=1}^{\infty} w_{i+1} \frac{\partial}{\partial w_i} \end{aligned}$$

It is easy to see that $D_x \left(\frac{w_3}{w_1} - \frac{3}{2} \left(\frac{w_2}{w_1} \right)^2 \right) = 0$ if and only if $a_{uuu} = 0$.

Thus we obtain the next

Theorem 2: *The differential substitution (1) satisfies our assumption if and only if it is equivalent to a substitution of the same form corresponding to the system (9) with $a = \alpha(x, v)u^2 + \beta(x, v)u + \gamma(x, v)$.*

To understand how many essentially different substitutions are described by the system (9) with $a_{uuu} = 0$, we need study the equivalence problem for the substitutions (1). It is convenient to do in terms of the system (9).

Transformations

$$\begin{aligned} x &= \alpha(\bar{x}) \\ u &= \varphi(\bar{x}, \bar{u}) \\ v &= \psi(\bar{x}, \bar{v}) \end{aligned} \quad (10)$$

of the system (9) preserve the form of the latter. These transformations correspond to the point transformations of the equations (2) and (3) which preserve the

form of the substitution (1). Contact transformations of the equations (2) and (3) preserving the form of (1) induce a transformation of the system (9)

$$\begin{aligned} x &= \varphi(\bar{x}, \bar{v}), \varphi_v \neq 0 \\ u &= \psi(\bar{x}, \bar{u}, \bar{v}) \\ v &= \Phi(\bar{x}, \bar{v}), \end{aligned} \quad (11)$$

where φ, ψ and Φ satisfy the condition

$$\psi_v - a(\varphi, \psi, \Phi)\varphi_v = 0 \quad (12)$$

The transformation (11)-(12) preserves the form of (9) and the right-hand side $\bar{a}$ of the transformed system is expressed by the formula

$$\bar{a} = \frac{\varphi_x \psi_v - \varphi_v \psi_x}{\varphi_v \psi_u}. \quad (13)$$

Using the formulas (12) and (13) we can prove the next proposition by a simple but tedious argument.

Proposition 2: *Let a transformation (10)-(13) relates a system*

$$\begin{cases} u_x = a(x, u, v) = \alpha(x, v)u^2 + \beta(x, v)u + \gamma(x, v) \\ v_y = 0 \end{cases},$$

where $(a_{uv}/a_v)_v \neq 0$, to a system

$$\begin{cases} \bar{u}_{\bar{x}} = \bar{a}(\bar{x}, \bar{u}, \bar{v}) = \bar{\alpha}(\bar{x}, \bar{v})\bar{u}^2 + \bar{\beta}(\bar{x}, \bar{v})\bar{u} + \bar{\gamma}(\bar{x}, \bar{v}) \\ \bar{v}_y = 0 \end{cases},$$

where $(\bar{a}_{\bar{u}\bar{v}}/\bar{a}_{\bar{v}})_{\bar{v}} \neq 0$. Then this transformation must be either of the form

$$\begin{aligned} x &= \varphi(\bar{x}, \bar{v}) \\ v &= \Phi(\bar{x}, \bar{v}) \\ u &= \frac{\lambda(\bar{x}, \bar{v})}{\bar{u} + \eta(\bar{x}, \bar{v})} + \xi(\bar{x}, \bar{v}), \end{aligned} \quad (14)$$

where λ, η and ξ satisfy the conditions

$$\begin{aligned} \eta_{\bar{v}} &= -\lambda\varphi_{\bar{v}}\alpha(\varphi, \Phi) \\ \eta_{\bar{x}} &= \eta\bar{\beta} - \eta^2\bar{\alpha} - \bar{\gamma} - \lambda\varphi_{\bar{x}}\alpha(\varphi, \Phi) \\ \lambda_{\bar{v}} &= \lambda\varphi_{\bar{v}}(\beta(\varphi, \Phi) + 2\xi\alpha(\varphi, \Phi)) \\ \lambda_{\bar{x}} &= \lambda(\varphi_{\bar{x}}(2\alpha(\varphi, \Phi)\xi + \beta(\varphi, \Phi)) + \bar{\beta} - 2\eta\bar{\alpha}) \\ \xi_{\bar{v}} &= \varphi_{\bar{v}}(\alpha(\varphi, \Phi)\xi^2 + \beta(\varphi, \Phi)\xi + \gamma(\varphi, \Phi)) \\ \xi_{\bar{x}} &= \varphi_{\bar{x}}(\alpha(\varphi, \Phi)\xi^2 + \beta(\varphi, \Phi)\xi + \gamma(\varphi, \Phi)) + \lambda\bar{\alpha}, \end{aligned}$$

or of the form

$$\begin{aligned} x &= \varphi(\overline{x}) \\ v &= \Phi(\overline{x}, \overline{v}) \\ u &= \lambda(\overline{x})\overline{u} + \xi(\overline{x}). \end{aligned} \tag{15}$$

In both cases $\lambda \neq 0$.

We need the above proposition to prove the following

Proposition 3: *The differential substitutions described by the system (9) with $a = u^2 + uv + v^\gamma$, where γ is constant, $\gamma \neq 0, 1$, are pairwise nonequivalent for different γ .*

To prove this proposition it is sufficient to verify that the system

$$\begin{aligned} \eta_{\overline{v}} &= -\lambda\varphi_{\overline{v}} \\ \eta_{\overline{x}} &= \eta_{\overline{v}} - \eta^2 - \overline{v}^r - \lambda\varphi_{\overline{x}} \\ \lambda_{\overline{v}} &= \lambda\varphi_{\overline{v}}(\Phi + 2\xi) \\ \lambda_{\overline{x}} &= \lambda(\varphi_{\overline{x}}(\Phi + 2\xi) + \overline{v} - 2\eta) \\ \xi_{\overline{v}} &= \varphi_{\overline{v}}(\xi^2 + \Phi\xi + \Phi^s) \\ \xi_{\overline{x}} &= \varphi_{\overline{x}}(\xi^2 + \Phi\xi + \Phi^s) + \lambda \end{aligned}$$

is incompatible if $r \neq s$. Then it should be checked that the differential substitutions of the above form are nonequivalent under point transformations (15).

In conclusion, we give without a proof the next results which are obtained by the more detailed study of the equivalence problem for the substitutions (1).

Proposition 4: *The differential substitution (1) is equivalent to a differential substitutions of the form $v = u_x + p(x, u)$ if and only if the right-hand side a of the corresponding system (9) satisfy one of the following conditions:*

$$\left(\frac{a_{vv}}{a_v} \right)_u = 0 \quad \text{or} \quad \left(\frac{\left(\frac{a_{xv} + a a_{uv}}{a_v} - a_u \right)_u}{\left(\frac{a_{vv}}{a_v} \right)_u} \right)_u = 0$$

Theorem 3: *Any differential substitution (1) described by the system (9) with $a_{uuu} = 0$ is equivalent to one of the following substitutions:*

A) $v = u_x$

B) $v = u_x + e^u$

C) $v = u_x + e^u + e^{-u}$

D) $v = u_x + u^2$

F) *the substitutions described by the system (9) with $a = u^2 + vu + \gamma(x, v)$, $\gamma_{vv} \neq$*

0.

The cases A, B, C, D and F are nonequivalent.

Using proposition 4 we easily verify that the case F can not be reduced to the form $v = u_x + p(x, u)$. Thus there exist four differential substitutions of the form $v = u_x + p(x, u)$ and the infinite set of substitutions which are not equivalent to any substitution $v = u_x + p(x, u)$.

Acknowledgments

The author wishes to thank S. I. Svinolupov for many useful discussions. The author especially thanks V. V. Sokolov who offered the idea of this research and recommended to apply the technique of the generalized Laplace transformations to the present work. The idea of the proof of proposition 1 also belongs to V. V. Sokolov.

References.

- [1] V. V. Sokolov. *On the symmetries of evolutions equations*. Russian Math. Surv., 43(5), 1988, 165-204.
- [2] I. M. Anderson and N. Kamran. *The variational bicomplex for second order scalar partial differential equations in the plane*. CMR technical report, September, 1994.
- [3] V. V. Sokolov and A. V. Zhiber. *On the Darboux integrable hyperbolic equations*. To appear in Phis. Lett. A.
- [4] A. V. Zhiber, V. V. Sokolov and S. Ya. Startsev. *On the Darboux integrable nonlinear hyperbolic equations*. To appear in Doklady RAN (in Russian).
- [5] M. Juras and I.M Anderson. *Generalized Laplace invariant and the Method of Darboux*. Preprint, University of Montreal, June, 1995.
- [6] A. V. Zhiber. *Quasilinear hyperbolic equations with an infinite-dimensional algebra of symmetries*. Doclady RAN, v.58, No 4, 1994 (in Russian).